

Once More about Spectral-Dependent Quantum R -Matrix for $U_q(A_2)$

Anthony J. Bracken, Mark D. Gould and Yao-Zhong Zhang

Department of Mathematics, University of Queensland, Brisbane, Qld 4072, Australia

Abstract:

The recently obtained results in [1] are used to compute the explicitly spectral-dependent quantum R -matrix (or the intertwiners) on $V_{(6)}(x) \otimes V_{(6)}(y)$ and $V_{(3)}(x) \otimes V_{(6)}(y)$, where $V_{(6)}$ and $V_{(3)}$ are the 6-dimensional and fundamental representations of $U_q(A_2)$, respectively. It appears that the R -matrix on $V_{(3)}(x) \otimes V_{(6)}(y)$ depends on q in the different way from what one might usually think: q occurs in the R -matrix in fractional powers. It seems to be the first example in literatures of R -matrix with the new feature.

PACS numbers: 03.65.Fd ; 02.20.+b

In our previous papers [1] we determine the explicitly spectral-dependent universal R -matrix for the quantum algebras $U_q(A_1)$ and $U_q(A_2)$ by using the universal R -matrix [2][3] for the corresponding quantum affine algebras $U_q(A_1^{(1)})$ and $U_q(A_2^{(1)})$, respectively; applying the results to the adjoint representation of $U_q(A_2)$, the simplest non-trivial case where the tensor product of the representation with itself is with finite multiplicity, we obtain the explicitly spectral-dependent solution to the quantum Yang-Baxter equation [4][5][6] associated to the the adjoint representation of $U_q(A_2)$.

The present letter continues the above investigations to other interesting cases to find the spectral-dependent quantum R -matrix on modules $V_{(6)}(x) \otimes V_{(6)}(y)$ and $V_{(3)}(x) \otimes V_{(6)}(y)$, where $V_{(6)}$ and $V_{(3)}$ are the 6-dimensional and fundamental representations of $U_q(A_2)$, respectively.

The former may be in principal treated by means of the usual "Yang-Baxterization" procedure used in literatures [7][5][8][9], however, trying to find an explicit expression for the R -matrix by this procedure is inevitably involved in the determination of the explicit CG coefficients for $U_q(A_2)$ which as is well-known is very complicated and is not generally available even for the corresponding classical algebra A_2 . The latter could not be treated by the "Yang-Baxterization" approach since braid group generators always act on the tensored space of the *same* representation. This may explain why all previous attempts in literatures at finding spectral-dependent R -matrix are only limited in the case where the representations being tensored are same. On the contrast, our approach developed in [1] not only gives rise to an unified treatment for the case of whether the tensor product is multiplicity-free or with finite multiplicity, as showed in [1], but also to an unified treatment for the case of whether the representations being tensored are same or different, as we will see below.

There is more to that. As one may see in (17) below, the R -matrix or the intertwiner on $V_{(3)}(x) \otimes V_{(6)}(y)$ behaves different from what one might usually think: q appears in the R -matrix in the form of its fractional powers. To our knowledge, it is the first example in literatures of quantum R -matrix with the new feature. Therefore, the present study should throw light on new kind of solutions to the quantum Yang-Baxter equation.

We begin with a brief review on some results obtained in [1]. Throughout this letter we use

the notations

$$\begin{aligned} (n)_q &= \frac{1-q^n}{1-q}, \quad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \quad q_\alpha = q^{(\alpha, \alpha)} \\ \exp_q(x) &= \sum_{n \geq 0} \frac{x^n}{(n)_q!}, \quad (n)_q! = (n)_q(n-1)_q \dots (1)_q \end{aligned} \quad (1)$$

In ref.[1], starting from the universal R -matrix for the quantum (non-twisted) affine algebra $U_q(A_2^{(1)})$, we determine the explicitly spectral-dependent universal R -matrix $R(x, y)$ for the corresponding algebra $U_q(A_2)$, given by,

$$\begin{aligned} R(x, y) &= \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n \left(q^{-nh_\alpha} E_\alpha q^{-n(h_\alpha + 2h_\beta)/3} \otimes q^{n(h_\alpha + 2h_\beta)/3} F_\alpha q^{nh_\alpha} \right) \right) \\ &\cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n \left(q^{-nh_{\alpha+\beta}} E_{\alpha+\beta} q^{n(h_\beta - h_\alpha)/3} \otimes q^{n(h_\alpha - \beta)/3} F_{\alpha+\beta} q^{nh_{\alpha+\beta}} \right) \right) \\ &\cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n \left(E'_{\beta+n\delta} \otimes F'_{\beta+n\delta} \right) \right) \\ &\cdot \exp \left(\sum_{n > 0} \sum_{i, j=1}^2 C_{ij}^n(q) (q - q^{-1}) \left(\frac{x}{y} \right)^n (E'_{n\delta}^{(i)} \otimes F'_{n\delta}^{(j)}) \right) \\ &\cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(E'_{(\delta-\beta)+n\delta} \otimes F'_{(\delta-\beta)+n\delta} \right) \right) \\ &\cdot \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(q^{-(n+1)(h_\alpha + 2h_\beta)/3} F_\alpha q^{-nh_\alpha} \otimes q^{nh_\alpha} E_\alpha q^{(n+1)(h_\alpha + 2h_\beta)/3} \right) \right) \\ &\cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(q^{(n+1)(h_\beta - h_\alpha)/3} F_{\alpha+\beta} q^{-nh_{\alpha+\beta}} \right. \right. \\ &\quad \left. \left. \otimes q^{nh_{\alpha+\beta}} E_{\alpha+\beta} q^{(n+1)(h_\alpha - h_\beta)/3} \right) \right) \cdot q^{\sum_{i, j=1}^2 (a_{\text{sym}}^{-1})^{ij} h_i \otimes h_j} \end{aligned} \quad (2)$$

where α and β are the simple roots of A_2 and

$$\begin{aligned} E_{\alpha+\beta} &= E_\alpha E_\beta - q^{-1} E_\beta E_\alpha, \quad F_{\alpha+\beta} = F_\beta F_\alpha - q F_\alpha F_\beta \\ (a_{\text{sym}}^{ij}) &= \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \\ (C_{ij}^m(q)) &= (C_{ji}^m(q)) = \frac{n}{[n]_q} \frac{[2]_q^2}{q^{2n} + 1 + q^{-2n}} \begin{pmatrix} q^n + q^{-n} & (-1)^n \\ (-1)^n & q^n + q^{-n} \end{pmatrix} \\ E'_{\beta+n\delta} &= (-1)^n [2]_q^{-n} q^n \left\{ (\text{ad}'_{q^{-1}} \mathcal{E})^n E_\beta \right\} q^{n(h_\beta - h_\alpha)/3} \\ F'_{\beta+n\delta} &= [2]_q^{-n} q^{n(h_\alpha - h_\beta)/3} (\text{ad}'_{q^{-1}} \mathcal{F})^n F_\beta \\ E'_{(\delta-\beta)+n\delta} &= [2]_q^{-n} q^{-n} \left\{ (\text{ad}'_q \mathcal{E})^n (\text{ad}'_{q^{-2}} E_\alpha) F_{\alpha+\beta} \right\} q^{(n+1)(h_\beta - h_\alpha)/3} \\ F'_{(\delta-\beta)+n\delta} &= (-1)^n [2]_q^{-n} q^{(n+1)(h_\alpha - h_\beta)/3} (\text{ad}'_q \mathcal{F})^n (\text{ad}'_{q^2} E_{\alpha+\beta}) F_\alpha \end{aligned}$$

$$\mathcal{E} = (\text{ad}'_{q^{-1}} E_\beta)(\text{ad}'_{q^{-2}} E_\alpha) F_{\alpha+\beta}, \quad \mathcal{F} = (\text{ad}'_q (\text{ad}'_{q^2} E_{\alpha+\beta}) F_\alpha) F_\beta, \quad (3)$$

$(\text{ad}'_Q \mathcal{A}) \cdot \mathcal{B} \equiv \mathcal{A}\mathcal{B} - Q\mathcal{B}\mathcal{A}$; $E'_{n\delta}{}^{(i)}$ and $F'_{n\delta}{}^{(i)}$ are determined by the equalities of formal series:

$(\alpha_i = \alpha, \beta)$

$$\begin{aligned} (q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{k=1}^{\infty} \tilde{E}'_{k\delta}{}^{(i)} u^k &= \exp \left((q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{l=1}^{\infty} \tilde{E}'_{l\delta}{}^{(i)} u^l \right) - 1 \\ -(q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{k=1}^{\infty} \tilde{F}'_{k\delta}{}^{(i)} u^{-k} &= \exp \left(-(q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{l=1}^{\infty} \tilde{F}'_{l\delta}{}^{(i)} u^{-l} \right) - 1 \end{aligned} \quad (4)$$

in which

$$\begin{aligned} \tilde{E}'_{n\delta}{}^{(\alpha)} &= (-1)^{n-1} [2]_q^{-1} (E_\alpha F_\alpha - q^{-2n} F_\alpha E_\alpha) q^{-(n-1)h_\alpha} q^{-n(h_\alpha+2h_\beta)/3} \\ \tilde{F}'_{n\delta}{}^{(\alpha)} &= (-1)^{n-1} [2]_q^{-1} q^{(n-1)h_\alpha} q^{n(h_\alpha+2h_\beta)/3} (F_\alpha E_\alpha - q^{2n} E_\alpha F_\alpha) \\ \tilde{E}'_{n\delta}{}^{(\beta)} &= (-1)^n [2]_q^{-n} q^{n-2} \left\{ (\text{ad}'_{q^{-n+2}} \mathcal{F}') \cdot (\text{ad}'_{q^{-1}} \mathcal{E})^{n-1} E_\beta \right\} q^{n(h_\beta-h_\alpha)/3} \\ \tilde{F}'_{n\delta}{}^{(\beta)} &= [2]_q^{-n} q^{n-1} q^{n(h_\alpha-h_\beta)/3} (\text{ad}'_{q^{-n+2}} \mathcal{E}') \cdot (\text{ad}'_{q^{-1}} \mathcal{F})^{n-1} F_\beta \\ \mathcal{E}' &= E_{\alpha+\beta} F_\alpha - q^2 F_\alpha E_{\alpha+\beta}, \quad \mathcal{F}' = E_\alpha F_{\alpha+\beta} - q^{-2} F_{\alpha+\beta} E_\alpha \end{aligned} \quad (5)$$

It is shown in our previous papers [1] that the explicit form of generators on the fundamental representation of $U_q(A_2)$ is given by

$$\begin{aligned} h_\alpha &= \text{diag}(1, -1, 0), \quad h_\beta = \text{diag}(0, 1, -1) \\ E_\alpha &= e_{12}, \quad F_\alpha = e_{21}, \quad E_{\alpha+\beta} = e_{13}, \quad F_{\alpha+\beta} = e_{31} \\ E'_{\beta+n\delta} &= q^{-2n-n/3} e_{23}, \quad F'_{\beta+n\delta} = q^{2n+n/3} e_{32} \\ E'_{(\delta-\beta)+n\delta} &= -q^{-2n-1-(n+1)/3} e_{32}, \quad F'_{(\delta-\beta)+n\delta} = -q^{2n+1+(n+1)/3} e_{23} \\ E'_{n\delta}{}^{(\alpha)} &= [2]_q^{-1} (-1)^{n-1} \frac{[n]_q}{n} q^{-n/3} (e_{11} - q^{-2n} e_{22}) \\ F'_{n\delta}{}^{(\alpha)} &= [2]_q^{-1} (-1)^{n-1} \frac{[n]_q}{n} q^{n/3} (e_{11} - q^{2n} e_{22}) \\ E'_{n\delta}{}^{(\beta)} &= -[2]_q^{-1} \frac{[n]_q}{n} q^{-n/3} (q^{-n} e_{22} - q^{-3n} e_{33}) \\ F'_{n\delta}{}^{(\beta)} &= -[2]_q^{-1} \frac{[n]_q}{n} q^{n/3} (q^n e_{22} - q^{3n} e_{33}) \end{aligned} \quad (6)$$

where (and below) e_{ij} is the matrix satisfying $(e_{ij})_{kl} = \delta_{ik}\delta_{jl}$ and $e_{ij}e_{kl} = \delta_{jk}e_{il}$. Upon applying

(2) to $V_{(3)} \otimes V_{(3)}$ in which $V_{(3)}$ denotes the fundamental representation of $U_q(A_2)$, we get [1]

$$\begin{aligned} R_{(3),(3)}(x, y) &= f_q(x, y) \cdot \left(e_{11} + e_{99} + \frac{q^{-1}(y-x)}{y-q^{-2}x} (e_{22} + e_{33} + e_{44} + e_{66} + e_{77} + e_{88}) + \right. \\ &\quad \left. + \frac{y(1-q^{-2})}{y-q^{-2}x} (e_{24} + e_{37} + e_{68}) + \frac{x(1-q^{-2})}{y-q^{-2}x} (e_{42} + e_{73} + e_{86}) \right) \end{aligned} \quad (7)$$

where

$$f_q(x, y) = q^{2/3} \cdot \exp \left(\sum_{n>0} \frac{q^{2n} - q^{-2n}}{q^{2n} + 1 + q^{-2n}} \frac{(x/y)^n}{n} \right) \quad (8)$$

Eq.(7) is the well known result, up to the scalar factor $f_q(x, y)$, due to Jimbo [5]. We reproduce it by our approach.

We now come to our main concern in this letter. First we want to apply (2) to $V_{(6)} \otimes V_{(6)}$, to find the quantum R -matrix, $R_{(6),(6)}(x, y)$, on the tensor product module for $U_q(A_2)$. As we did in [1], we introduce the so-called Gelfand-Tsetlin basis vector $|(m) \rangle$ given by

$$|(m) \rangle = \left| \begin{pmatrix} m_{13} & m_{23} & m_{33} \\ & m_{12} & m_{22} \\ & & m_{11} \end{pmatrix} \right\rangle \quad (9)$$

For the 6-dimensional representation, we have the following 6 basis vectors:

$$\begin{aligned} \phi_1 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & 4/3 & -2/3 \\ & & 4/3 \end{pmatrix} \right\rangle, & \phi_2 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & 4/3 & -2/3 \\ & & 1/3 \end{pmatrix} \right\rangle \\ \phi_3 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & 4/3 & -2/3 \\ & & -2/3 \end{pmatrix} \right\rangle, & \phi_4 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & 1/3 & -2/3 \\ & & 1/3 \end{pmatrix} \right\rangle \\ \phi_5 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & 1/3 & -2/3 \\ & & -2/3 \end{pmatrix} \right\rangle, & \phi_6 &= \left| \begin{pmatrix} 4/3 & -2/3 & -2/3 \\ & -2/3 & -2/3 \\ & & -2/3 \end{pmatrix} \right\rangle \end{aligned} \quad (10)$$

Then long computations plus induction in n show that the matrix form of generators in the 6-dimensional representation of $U_q(A_2)$ is given by

$$\begin{aligned} h_\alpha &= \text{diag}(2, 0, -2, 1, -1, 0), & h_\beta &= \text{diag}(0, 1, 2, -1, 0, -2) \\ E_\alpha &= [2]_q^{1/2} e_{12} + [2]_q^{1/2} e_{23} + e_{45}, & F_\alpha &= [2]_q^{1/2} e_{21} + [2]_q^{1/2} e_{32} + e_{54} \\ E_\beta &= e_{24} + [2]_q^{1/2} e_{35} + [2]_q^{1/2} e_{56}, & F_\beta &= e_{42} + [2]_q^{1/2} e_{53} + [2]_q^{1/2} e_{65} \\ E_{\alpha+\beta} &= [2]_q^{1/2} e_{14} + q e_{25} + [2]_q^{1/2} e_{46}, & F_{\alpha+\beta} &= [2]_q^{1/2} e_{41} + q e_{52} + [2]_q^{1/2} e_{64} \\ E'_{\beta+n\delta} &= q^{-2n/3} \{ q^{-3n} e_{24} + [2]_q^{1/2} q^{-3n} e_{35} + [2]_q^{1/2} q^{-n} e_{56} \} \end{aligned}$$

$$\begin{aligned}
F'_{\beta+n\delta} &= q^{2n/3} \{q^{3n} e_{42} + [2]_q^{1/2} q^{3n} e_{53} + [2]_q^{1/2} q^n e_{65}\} \\
E'_{(\delta-\beta)+n\delta} &= -q^{-2(n+1)/3} \{q^{-3n-2} e_{42} + [2]_q^{1/2} q^{-3n-1} e_{53} + [2]_q^{1/2} q^{-n-1} e_{65}\} \\
F'_{(\delta-\beta)+n\delta} &= -q^{2(n+1)/3} \{q^{3n+2} e_{24} + [2]_q^{1/2} q^{3n+1} e_{35} + [2]_q^{1/2} q^{n+1} e_{56}\} \\
E'_{n\delta}^{(\alpha)} &= [2]_q^{-1} (-1)^{n-1} \frac{[n]_q}{n} q^{-2n/3} \left\{ (q^n + q^{-n}) e_{11} + q^{-n} (q^{2n} - q^{-2n}) e_{22} \right. \\
&\quad \left. - q^{-2n} (q^n + q^{-n}) e_{33} + q^n e_{44} - q^{-n} e_{55} \right\} \\
F'_{n\delta}^{(\alpha)} &= [2]_q^{-1} (-1)^{n-1} \frac{[n]_q}{n} q^{2n/3} \left\{ (q^n + q^{-n}) e_{11} - q^n (q^{2n} - q^{-2n}) e_{22} \right. \\
&\quad \left. - q^{2n} (q^n + q^{-n}) e_{33} + q^{-n} e_{44} - q^n e_{55} \right\} \\
E'_{n\delta}^{(\beta)} &= [2]_q^{-1} \frac{[n]_q}{n} q^{-2n/3} \left\{ -q^{-2n} e_{22} - q^{-n} (q^n + q^{-n}) e_{33} + q^{-4n} e_{44} \right. \\
&\quad \left. - q^{-2n} (q^{2n} - q^{-2n}) e_{55} + q^{-3n} (q^n + q^{-n}) e_{66} \right\} \\
F'_{n\delta}^{(\beta)} &= -[2]_q^{-1} \frac{[n]_q}{n} q^{2n/3} \left\{ -q^{2n} e_{22} - q^n (q^n + q^{-n}) e_{33} + q^{4n} e_{44} \right. \\
&\quad \left. + q^{2n} (q^{2n} - q^{-2n}) e_{55} + q^{3n} (q^n + q^{-n}) e_{66} \right\}
\end{aligned} \tag{11}$$

We have the following properties for the generators in (11),

$$\begin{aligned}
(E_\alpha)^2 &= [2]_q e_{13}, \quad (E_\alpha)^3 = 0, \quad (F_\alpha)^2 = [2]_q e_{31}, \quad (F_\alpha)^3 = 0 \\
(E_{\alpha+\beta})^2 &= -[2]_q e_{16}, \quad (E_{\alpha+\beta})^3 = 0, \quad (F_{\alpha+\beta})^2 = -[2]_q e_{61}, \quad (F_{\alpha+\beta})^3 = 0 \\
(E'_{\beta+n\delta})^2 &= [2]_q q^{-5n-n/3} e_{36}, \quad (E'_{\beta+n\delta})^3 = 0, \quad (F'_{\beta+n\delta})^2 = [2]_q q^{5n+n/3} e_{63}, \quad (F'_{\beta+n\delta})^3 = 0 \\
(E'_{(\delta-\beta)+n\delta})^2 &= [2]_q q^{-5n-3-(n+1)/3} e_{63}, \quad (E'_{(\delta-\beta)+n\delta})^3 = 0 \\
(F'_{(\delta-\beta)+n\delta})^2 &= [2]_q q^{5n+3+(n+1)/3} e_{36}, \quad (F'_{(\delta-\beta)+n\delta})^3 = 0
\end{aligned} \tag{12}$$

as can be easily checked.

Inserting (11) into (2), we see that in the expansion of each q -exponential only three terms survive thanks to the celebrated properties of generators, eq.(12). Thus one is able to work out the infinite products in (2). The contributions from the imaginary root vectors in (2) can also be worked out term by term and written as a very compact form. The final result may be put in the explicit and compact form,

$$\begin{aligned}
R_{(6),(6)}(x, y) &= \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n (E'_{\alpha+n\delta} \otimes F'_{\alpha+n\delta}) + y^2 f(e_{13} \otimes e_{31}) \right\} \\
&\quad \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n (E'_{\alpha+\beta+n\delta} \otimes F'_{\alpha+\beta+n\delta}) + y^2 f(e_{16} \otimes e_{61}) \right\}
\end{aligned}$$

$$\begin{aligned}
& \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n \left(E'_{\beta+n\delta} \otimes F'_{\beta+n\delta} \right) + y^2 f(e_{36} \otimes e_{63}) \right\} \\
& \cdot q^{-1} f_q(x, y) \cdot \{\text{imaginary root vectors contribution}\} \\
& \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} \left(E'_{(\delta-\beta)+n\delta} \otimes F'_{(\delta-\beta)+n\delta} \right) + x^2 f(e_{63} \otimes e_{36}) \right\} \\
& \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} \left(E'_{(\delta-\alpha)+n\delta} \otimes F'_{(\delta-\alpha)+n\delta} \right) + x^2 f(e_{31} \otimes e_{13}) \right\} \\
& \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} \left(E'_{(\delta-\alpha-\beta)+n\delta} \otimes F'_{(\delta-\alpha-\beta)+n\delta} \right) + x^2 f(e_{61} \otimes e_{16}) \right\} \\
& \cdot \left\{ q^3(e_{11} \otimes e_{11} + e_{33} \otimes e_{33} + e_{66} \otimes e_{66}) + q(e_{22} \otimes e_{22} + \right. \\
& + e_{44} \otimes e_{44} + e_{55} \otimes e_{55}) + q(e_{11} \otimes e_{22} + e_{11} \otimes e_{44} + e_{22} \otimes e_{33} + \\
& + e_{33} \otimes e_{55} + e_{44} \otimes e_{66} + e_{55} \otimes e_{66} + \{\leftrightarrow\}) + q^{-1}(e_{11} \otimes e_{33} + e_{11} \otimes e_{55} + \\
& + e_{11} \otimes e_{66} + e_{22} \otimes e_{66} + e_{33} \otimes e_{44} + e_{33} \otimes e_{66} + \{\leftrightarrow\}) + \\
& \left. + (e_{22} \otimes e_{44} + e_{22} \otimes e_{55} + e_{44} \otimes e_{55} + \{\leftrightarrow\}) \right\} \tag{13}
\end{aligned}$$

where $f_q(x, y)$ is given by (8) and " $\{\leftrightarrow\}$ " denotes the interchange of the quantities in the space

$X \otimes Y$; $E'_{\beta+n\delta}$, $F'_{\beta+n\delta}$, $E'_{(\delta-\beta)+n\delta}$, $F'_{(\delta-\beta)+n\delta}$ are given in (11) and

$$\begin{aligned}
f &= [2]_q q^{-1} (q - q^{-1})^2 \frac{y + q^4 x}{(y^2 - x^2)(y - q^2 x)} \\
E'_{\alpha+n\delta} &= (-1)^n q^{-2n/3} \left\{ [2]_q^{1/2} q^{-2n} e_{12} + [2]_q^{1/2} e_{23} + e_{45} \right\} \\
F'_{\alpha+n\delta} &= (-1)^n q^{2n/3} \left\{ [2]_q^{1/2} q^{2n} e_{21} + [2]_q^{1/2} e_{32} + e_{54} \right\} \\
E'_{\alpha+\beta+n\delta} &= (-1)^n q^{-2n/3} \left\{ [2]_q^{1/2} q^{-2n} e_{14} + q e_{25} + [2]_q^{1/2} e_{46} \right\} \\
F'_{\alpha+\beta+n\delta} &= (-1)^n q^{2n/3} \left\{ [2]_q^{1/2} q^{2n} e_{41} + q e_{52} + [2]_q^{1/2} e_{64} \right\} \\
E'_{(\delta-\alpha)+n\delta} &= (-1)^n q^{-2(n+1)/3} \left\{ [2]_q^{1/2} q^{-2n} e_{21} + [2]_q^{1/2} e_{32} + e_{54} \right\} \\
F'_{(\delta-\alpha)+n\delta} &= (-1)^n q^{2(n+1)/3} \left\{ [2]_q^{1/2} q^{2n} e_{12} + [2]_q^{1/2} e_{23} + e_{45} \right\} \\
E'_{(\delta-\alpha-\beta)+n\delta} &= (-1)^n q^{-2(n+1)/3} \left\{ [2]_q^{1/2} q^{-2n} e_{41} + e_{52} + [2]_q^{1/2} e_{64} \right\} \\
F'_{(\delta-\alpha-\beta)+n\delta} &= (-1)^n q^{2(n+1)/3} \left\{ [2]_q^{1/2} q^{2n} e_{14} + e_{25} + [2]_q^{1/2} e_{46} \right\} \\
\{\text{imaginary root vectors contribution}\} &= a'/a \sum_{i=1}^6 (1 + (b/b' - 1)(\delta_{i2} + \delta_{i4} + \delta_{i5})) (e_{ii} \otimes e_{ii}) \\
&+ ba'(e_{11} \otimes e_{22} + e_{11} \otimes e_{44}) + ab(e_{11} \otimes e_{33} + e_{11} \otimes e_{55} + e_{11} \otimes e_{66}) + \\
&+ 1/(ab')(e_{22} \otimes e_{11}) + a'b(e_{22} \otimes e_{33}) + ba'/b'(e_{22} \otimes e_{44}) + aa'b(e_{22} \otimes e_{55}) + \\
&+ ab(e_{22} \otimes e_{66}) + 1/(a'b')(e_{33} \otimes e_{11}) + 1/(ab')(e_{33} \otimes e_{22}) + b/b'(e_{33} \otimes e_{44}) +
\end{aligned}$$

$$\begin{aligned}
& + ba'(e_{33} \otimes e_{55}) + ab(e_{33} \otimes e_{66}) + 1/(ab')(e_{44} \otimes e_{11}) + b/(ab')(e_{44} \otimes e_{22}) + \\
& + b/b'(e_{44} \otimes e_{33}) + b/(a'b')(e_{44} \otimes e_{55}) + a'b(e_{44} \otimes e_{66}) + 1/(a'b')(e_{55} \otimes e_{11}) + \\
& + 1/(aa'b')(e_{55} \otimes e_{22}) + 1/(ab')(e_{55} \otimes e_{33}) + b/(ab')(e_{55} \otimes e_{44}) + ba'(e_{55} \otimes e_{66}) + \\
& + 1/(b'a')(e_{66} \otimes e_{11} + e_{66} \otimes e_{22} + e_{66} \otimes e_{33}) + 1/(ab')(e_{66} \otimes e_{44}) + \\
& + 1/(ab')(e_{66} \otimes e_{55})
\end{aligned} \tag{14}$$

in which we have defined

$$a = \frac{y - q^2x}{y - x}, \quad a' = \frac{y - q^{-2}x}{y - x}, \quad b = \frac{y - q^4x}{y - q^2x}, \quad b' = \frac{y - q^{-4}x}{y - q^{-2}x} \tag{15}$$

We see that (13) is an extremely explicit formula: the sums in (13) can be easily worked out.

But for compactness, we leave the sums in their present form.

We then apply (2) to another interesting case: $V_{(3)} \otimes V_{(6)}$, to derive the quantum R -matrix, $R_{(3),(6)}(x, y)$, on this tensor product module. As we addressed in the introduction, this case could not be treated by means of the usual Yang-Baxterization procedure simply because the representations being tensored are different. With the help of (6) and (11) and through long but similar calculations leading to (13) one finds

$$R_{(3),(6)}(x, y) = g_q(x, y) \cdot R_+ R_0 R_- \tag{16}$$

where

$$\begin{aligned}
g_q(x, y) &= q^{1/3} \cdot \exp \left(\sum_{n>0} \frac{q^n - q^{-n}}{q^{2n} + 1 + q^{-2n}} \frac{(q^{1/3}x/y)^n}{n} \right) \\
R_+ &= I + \frac{y(q - q^{-1})}{y - q^{4/3}x} \{ [2]_q^{1/2} (e_{12} \otimes e_{21} + e_{13} \otimes e_{41} + e_{23} \otimes e_{53}) + e_{23} \otimes e_{42} \} \\
&+ \frac{y(q - q^{-1})}{y - q^{-2/3}x} \{ e_{12} \otimes e_{54} + [2]_q^{1/2} (e_{12} \otimes e_{32} + e_{13} \otimes e_{64}) + q^{-1} e_{13} \otimes e_{52} + \\
&+ [2]_q^{1/2} e_{23} \otimes e_{65} \} + \frac{y^2(q - q^{-1})^2}{(y - q^{4/3}x)(y - q^{-2/3}x)} e_{13} \otimes e_{52} \\
R_0 &= \frac{y - q^{-2/3}x}{y - q^{4/3}x} \{ e_{11} \otimes e_{11} + e_{22} \otimes e_{33} + e_{33} \otimes e_{66} \} + \\
&\frac{(y - q^{10/3}x)(y - q^{-2/3}x)}{(y - q^{4/3}x)^2} \{ e_{11} \otimes e_{22} + e_{11} \otimes e_{44} + e_{22} \otimes e_{55} \} \\
&+ \frac{y - q^{10/3}x}{y - q^{4/3}x} \{ e_{11} \otimes e_{33} + e_{11} \otimes e_{55} + e_{11} \otimes e_{66} + e_{22} \otimes e_{66} \} \\
&+ \frac{y - q^{-2/3}x}{y - q^{-8/3}x} \{ e_{22} \otimes e_{11} + e_{33} \otimes e_{11} + e_{33} \otimes e_{22} + e_{33} \otimes e_{33} \}
\end{aligned}$$

$$\begin{aligned}
& + \frac{(y - q^{-2/3}x)^2}{(y - q^{4/3}x)(y - q^{-8/3}x)} \{e_{22} \otimes e_{22} + e_{33} \otimes e_{44} + e_{33} \otimes e_{55}\} \\
& + \frac{(y - q^{10/3}x)(y - q^{-2/3}x)^2}{(y - q^{4/3}x)^2(y - q^{-8/3}x)} e_{22} \otimes e_{44} \\
R_- = & q\{e_{11} \otimes e_{11} + e_{22} \otimes e_{33} + e_{33} \otimes e_{66}\} + e_{11} \otimes e_{22} + e_{11} \otimes e_{44} \\
& + e_{22} \otimes e_{22} + e_{22} \otimes e_{55} + e_{33} \otimes e_{44} + e_{33} \otimes e_{55} + q^{-1}\{e_{11} \otimes e_{33} + \\
& + e_{11} \otimes e_{55} + e_{11} \otimes e_{66} + e_{22} \otimes e_{11} + e_{22} \otimes e_{44} + \\
& + e_{22} \otimes e_{66} + e_{33} \otimes e_{11} + e_{33} \otimes e_{22} + e_{33} \otimes e_{33} + \\
& + \frac{xq^{1/3}(q - q^{-1})}{y - q^{4/3}x} \{[2]_q^{1/2}(e_{21} \otimes e_{12} + e_{31} \otimes e_{14} + e_{32} \otimes e_{35}) + e_{32} \otimes e_{24}\} \\
& + \frac{xq^{-2/3}(q - q^{-1})}{y - q^{-2/3}x} \{[2]_q^{1/2}(e_{21} \otimes e_{23} + e_{31} \otimes e_{46}) + e_{21} \otimes e_{45} + e_{31} \otimes e_{25} + \\
& + [2]_q^{1/2}e_{32} \otimes e_{56}\} + \frac{x^2q^{2/3}(q - q^{-1})^2}{(y - q^{4/3}x)(y - q^{-2/3}x)} e_{31} \otimes e_{25} \tag{17}
\end{aligned}$$

It is understood that I in R_+ is a 18×18 unit matrix.

To summarize, we have obtained the explicit and compact formulae for the quantum R -matrix (or the intertwiners) on $V_{(6)}(x) \otimes V_{(6)}(y)$ and $V_{(3)}(x) \otimes V_{(6)}(y)$ of $U_q(A_2)$. In the latter case, an interesting phenomenon appears: the powers of q are no longer integers rather fractional numbers! We believe that it is the first example in the literature of quantum R -matrix with the feature. The possibly physical meanings of this new phenomenon remain to be seen.

Acknowledgements: Y.Z.Z. would like to thank N.J.MacKay for e-mail comments and drawing his attention to ref.[9]. The financial support from Australian Research Council is gratefully acknowledged.

References

- [1] Y.-Z.Zhang and M.D.Gould, *Quantum Affine Algebra and Universal R-Matrix with Spectral Parameter: I, II*, The University of Queensland preprints UQMATH-93-05, UQMATH-93-06, hep-th/9307007, hep-th/9307008
- [2] S.M.Khoroshkin and U.N.Tolstoy, *Lett.Math.Phys.* **24** (1992) 231; *Funkz.Analyz. i ego Pril.* **26** (1992) 85
- [3] Y.-Z.Zhang and M.D.Gould, *On universal R-Matrix for quantized nontwisted rank 3 affine Lie algebras*, hep-th/9303095, *Lett.Math.Phys.* (in press); *Unitarity and complete reducibility of certain modules over quantized affine Lie algebras*, hep-th/9303096, *J.Math.Phys.* (in press)
- [4] V.G.Drinfeld, *Proc. ICM, Berkeley* **1** (1986) 798
- [5] M.Jimbo, *Lett.Math.Phys.* **10** (1985) 63, *ibid* **11** (1986) 247; *Commun.Math.Phys.* **102** (1986) 247
- [6] N.Reshetikhin, *Quantized universal enveloping algebras, the Yang-Baxter equation and invariants of links: I, II*, preprints LOMI E-4-87, E-17-87; A.N.Kirillov and N.Reshetikhin, *Representations of the algebra $U_q(sl(2))$, q -orthogonal polynomials and invariants of links*, preprint LOMI E-9-88
- [7] V.F.R.Jones, *Int.J.Mod.Phys.* **B4** (1990) 701
- [8] A.J.Bracken, M.D.Gould and R.B.Zhang, *Mod.Phys.Lett.* **A5** (1990) 831; R.B.Zhang, M.D.Gould and A.J.Bracken, *Nucl.Phys.* **354** (1991) 625
- [9] N.J.MacKay, *J.Phys. A: Math.Gen.* **24** (1991) 4017